

Yuri Brosgol, M.D.
Diplomate in Child and General Neurology
Certified by A.B.P.N.

Neurology Continuum, P.C.
30 West End Ave, Bld.1, Brooklyn, NY 11235
Tel. (718) 648-4567; Fax:(718) 648-0407
e-mail: ybrosgo@msn.com

INFORMATION SHEET
For the Use of CLONIDINE

Indications:

CLONIDINE is FDA approved for treatment of

- ADHD (attention deficit hyperactivity disorder) in children
- Autistic disorder in children
- Elevated high pressure;

CLONIDINE is used by physicians OFF LABEL (without FDA approval but based on literature evidence and clinical experience) for treatment of :

- Irritability, hyperactivity, insomnia in children
- Tics and Tourette's Syndrome
- Prophylaxis of migraines
- Severe neuropathic pain
- Opioid withdrawal syndrome
- Postmenopausal syndrome

CLONIDINE may cause serious side effects such as:

- Increased blood pressure if stopped suddenly from higher doses
- There have been reported sudden deaths in four children taking Clonidine and Methylphenidate together, however these patient had other serious predisposing conditions.

there are NON-SERIOUS reactions reported occasionally by the **CLONIDINE** users such as:

- Somnolence, dizziness, fatigue,
- Headaches, dry mouth, rash,
- nausea, constipation,
- legs cramps and swelling.